

INFOGRAPHIC

SPIDS – simplified predictive intubation difficulty score ☆




Vinícius Caldeira Quintão ^{a,*}, Cláudia Marquez Simões ^b, Maria José Carvalho Carmona ^a

^a Universidade de São Paulo, Hospital das Clínicas, Faculdade de Medicina, São Paulo, SP, Brazil

^b Hospital Sírio-Libanês, São Paulo, SP, Brazil

Available online 4 November 2020

SPIDS - simplified predictive intubation difficulty score 153 PATIENTS WHO UNDERWENT HEAD AND NECK SURGERIES



MMT, Modified Mallampati Test; RHNM_{max}, Maximum range of head and neck motion; TMD, thyromental distance; RHTMD, Ratio of height thyromental distance; NPV, negative predictive value.

DOI of original article:

<https://doi.org/10.1016/j.bjane.2020.09.007>

☆ Article reference: Selvi O, Kahraman ST, Tulgar S, Senturk O, Serifsoy TE, Thomas D, et al. Effectiveness of simplified predictive intubation difficulty score and thyromental height in head and neck surgeries: an observational study. Rev Bras Anesthesiol. 2020;70:595–604. <https://doi.org/10.1016/j.bjane.2020.09.007>

* Corresponding author.

E-mail: vinicius.quintao@hc.fm.usp.br (V.C. Quintão).

<https://doi.org/10.1016/j.bjane.2020.10.008>

© 2020 Sociedade Brasileira de Anestesiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).