

Journal Pre-proof

Debridement of axillary necrotizing fasciitis under anesthetic blocks of the serratus plane and supraclavicular brachial plexus: a case report

Leonardo Saraiva Guimarães de Oliveira, Renata de Andrade Chaves



PII: S0104-0014(21)00143-3

DOI: <https://doi.org/10.1016/j.bjane.2021.02.055>

Reference: BJANE 744119

To appear in: *Brazilian Journal of Anesthesiology (English edition)*

Received Date: 20 December 2020

Accepted Date: 27 February 2021

Please cite this article as: Oliveira LSGd, Chaves RdA, Debridement of axillary necrotizing fasciitis under anesthetic blocks of the serratus plane and supraclavicular brachial plexus: a case report, *Brazilian Journal of Anesthesiology (English edition)* (2021), doi: <https://doi.org/10.1016/j.bjane.2021.02.055>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

BJAN-D-20-00408_Case Report

Debridement of axillary necrotizing fasciitis under anesthetic blocks of the serratus plane and supraclavicular brachial plexus: a case report

Leonardo Saraiva Guimarães de Oliveira*, Renata de Andrade Chaves

Hospital Metropolitano Odilon Behrens, Prefeitura Municipal de Belo Horizonte, Departamento de Anestesiologia, Belo Horizonte, MG, Brazil

***Corresponding author.**

E-mail: saraivaleo@gmail.com (L.S.G. Oliveira).

ORCID ID:

Leonardo Saraiva Guimarães de Oliveira - <https://orcid.org/0000-0002-2467-921X>

Renata de Andrade Chaves - <https://orcid.org/0000-0002-6896-491X>

Received 20 December 2020; accepted 27 February 2021

Abstract

The regional techniques for axillary analgesia are well established. However, few studies have investigated surgical anesthesia. In this report, extensive debridement of axillary necrotizing fasciitis, including the posteromedial region of the right arm, performed under exclusive regional anesthesia in a patient with probable difficult airway is described. The procedure was performed under a Serratus Plane Block (SPB) and supraclavicular brachial plexus block, guided by ultrasound, and performed with venous sedation. We observed satisfactory anesthesia 15-minutes after the intervention and efficient intraoperative pain control within the following 24 hours. Surgical axilla anesthesia is feasible with the described blocks.

KEYWORDS: Axilla; Brachial plexus blocks; Case reports; Intermediate back muscles; Regional anesthesia; Thoracic wall

Introduction

Interfacial thoracic blocks, including Erector Spinae Plane (ESP), pectoral (PECS I and II) and Serratus Plane Blocks (SPB), are established analgesic techniques for thoracic, breast and axilla surgeries and are less invasive alternatives to Thoracic Epidural (TEB) and Thoracic Paravertebral Blocks (TPVB).[1-3] However, in the literature, there is no consensus on the performance of these blocks for anesthetic purposes, especially for axillary surgeries. The axilla is a region of intricate anatomy and innervation, which makes it challenging to induce surgical anesthesia in this compartment.[1]

In the present report, the combination of the SPB and supraclavicular brachial plexus block (BPB), guided by ultrasound, was proposed to address extensive axillary necrotizing fasciitis, which included the posteromedial region of the right arm, in a probable difficult airway scenario.

Case report

Voluntary and informed consent was provided by the patient for the inclusion of his data in this report. The patient was a 55-year-old male with a height of 1.75-m and weight of 82 kg; he had diabetes mellitus type II and took 500 mg of metformin/day. The patient was referred to the operating room for emergency surgical debridement with diffuse necrotizing fasciitis and multiple abscesses in the right axilla, which had been ongoing for 10-days due to a probable complication of suppurative hidradenitis (Fig. 1A). The patient had reported fasting for 8-hours. The physical examination revealed that he was conscious, eupneic, and hemodynamically stable. The airway examination revealed that the patient had a Mallampati score of IV, mouth opening of less than 4 cm, upper lip bite test class of II and thyromental distance of less than 6 cm. The axillary inspection showed extensive necrotizing fasciitis affecting the entire anatomical pyramid, extending to the edges of the pectoral muscles and the posteromedial face of the right arm. The laboratory tests showed leukocytosis of $15,000 \text{ cell.mm}^{-3}$ with deviation, PCR 259 mg.dL^{-1} , hyperglycemia of 280 mg.dL^{-1} (corrected to 170 mg.dL^{-1} with 10 IU of regular insulin) and arterial gasometry showing adequate peripheral perfusion. After standard monitoring, antibiotic therapy with 400 mg of ciprofloxacin and 600 mg of IV clindamycin was instituted. Sedation was started with 1 mcg.kg^{-1} dexmedetomidine in a 10-minute bolus and 1 mg/kg ketamine and was maintained with $0.7 \text{ mcg.kg}^{-1}.\text{h}^{-1}$ dexmedetomidine. Spontaneous ventilation was maintained with O_2 support by a nasal catheter at 2 L.min^{-1} . The blocks were guided by ultrasound (GE LOGIQ V2®; General Electric Company, Wauwatosa, WI) with a linear transducer (6–13 MHz). When performing the SPB, with the patient in supine position and arm abducted, the probe was slid distally and laterally (in sagittal orientation) from the middle portion of the clavicle until 4th and 5th ribs appeared. The probe was reoriented to the coronal plane and tilted backward, identifying the Latissimus Dorsi Muscle (LDM); then, the Serratus Anterior Muscle (SAM) was visualized deep in relation to the LDM and superficial to the ribs. A 50 mm needle (Locoplex® 50 mm, Vygon, Ecoen,

France) was inserted from the posterior to anterior direction in the interfascial plane of these muscles, and 20 mL of ropivacaine 0.5% was injected (Fig. 1B). In the BPB, the transducer was positioned in the transverse plane, proximal to the clavicle. The transducer was inclined caudally to visualize the thoracic structures (subclavian artery, 1st rib, pleura, and brachial plexus); the plexus was noted as a collection of hyperechoic oval structures posterior and superficial to the artery. The needle was introduced in the same plane, and 15 mL of ropivacaine 0.5% was injected (Fig. 1C). A loss of sensitivity was observed in the lateral thorax wall of T2–T6 of the entire axilla and lateral region of the pectorals as well as in the entire right upper limb during the pinprick test performed 15-minutes after the blocks. A fibrobronchoscope, videolaryngoscope and laryngeal mask n°5 were prepared in case airway rescue was needed. The procedure lasted 1-hour and 30-minutes, with satisfactory sensorial block and RASS-2 sedation scale scores. Adjuvant therapy with 2g of dipyron, 30 mg of ketorolac and 4 mg of ondansetron was administered. There were no pain complaints (score of 0 on the visual analogue scale) within 2 hours in the PACU. Within the following 24 hours, there were also no episodes of pain; 2g of dipyron was administered on a fixed schedule every 6 hours, and there was no need for opioids.

Discussion

It is well known that acute and chronic pain can develop after breast and axilla surgery, and regional anesthesia help prevent this. Both TPVB and TEB are traditional techniques for regional chest anesthesia, but they are elaborate and are associated with latent catastrophic complications.[1,2] With the evolution of ultrasound-guided blocks, new analgesic techniques have emerged for this purpose. Thoracic interfascial blocks such as the ESP, PECS I and II and SPB have appeared as safer and easier alternatives to perform.[1,3] However, while analgesia in these regions is satisfactory, procedures performed exclusively with peripheral nerve anesthesia are not widely performed. The axilla is a region of complex anatomy and innervation, and it is not simple to induce surgical anesthesia in this compartment (Table 1).[1]

Anesthetic blocks were chosen because of the difficult airway condition and to avoid general anesthesia. However, it is known that the choice for regional anesthesia is not always the guarantee of safety. In cases of block failure or complications, the difficult airway would need to be addressed suddenly, adding risks to the situation. Clinicians may choose to perform blocks in this scenario, but it is mandatory to prepare a plan for the airway.[4]

PVTB induces axillary analgesia by accessing ventral thoracic roots, blocking the intercostobrachial and intercostal nerves. However, for anesthetic purposes, theoretically, there is a gap in the coverage of innervation since the medial brachial cutaneous nerve and the medial and lateral

pectoral, long thoracic and thoracodorsal nerves originate from the brachial plexus.[1] By blocking similar innervations, the ESP also promotes good axillary analgesia. Nevertheless, the same limitations remain as with the PVTB. De Cassai et al. (2019) proposed the use of the SPB with brachial plexus selective nerve blocks to prevent these issues. The authors accessed the long thoracic, thoracodorsal, medial, and lateral pectoral nerves individually, inducing anesthesia in the region.[3]

Radiological and cadaveric dissection studies after SPBs have evidenced that solutions spread to the axilla and lateral thoracic wall, stimulating the reproduction of this block in a surgical context.[1,2] With the SPB, the accessed nerves are responsible for most of the sensitive innervation of the axilla (intercostobrachial and lateral cutaneous branches of intercostal), further the long thoracic and thoracodorsal nerves, responsible for axillary myotomes. All these nerves are located between the fascias of the SAM and LDM.[2] Sanllorrente-Sebastian et al. (2020) reported 2 cases of axillary arteriovenous fistulas being successfully treated under the supraclavicular BPB and SPB (between the SAM and external intercostal muscle at the 2nd rib level). In one case, it was necessary to additionally administer mepivacaine in the incision.[5] The ability to perform the technique in the supine position in patients with severe pain was the reason why the SPB was chosen over the ESP. The SPB was chosen over the PECS II (pectorals, long thoracic, intercostobrachial and lateral cutaneous branches of intercostal nerves are reached) because there was an infection that extended towards the anterosuperior chest. The reason that the BPB was performed was that the debridement extended to the posterior arm region (radial nerve territory) and there was a probable need to address the remaining axillary innervation (medial and lateral pectoral nerves and medial brachial cutaneous nerve).[1]

The SPB is a well-explored analgesic technique used in axillary surgeries. However, there is still a lack of strong evidence supporting the use of the SPB for anesthetic purposes in this context. In the present case, we successfully performed extensive axillary debridement with the SPB and supraclavicular BPB. Clinical trials are necessary to validate the presented conclusions as well as to determine the viability of the SPB as a unique technique in these circumstances and the required doses for this technique.

Authors' contributions

Leonardo Saraiva Guimarães de Oliveira: Conception and design of the study, data acquisition and interpretation, Writing of the article and critical review of it, Final approval of the version to be submitted.

Renata de Andrade Chaves: Conception and design of the study, data acquisition and interpretation, Critical review of the article, Final approval of the version to be submitted.

Conflicts of interest

The authors declare no conflicts of interest.

Glossary

BPB - brachial plexus block; ESP - erector spinae plane block; LDM - latissimus dorsi muscle; PECS - pectoralis blocks; RASS - Richmond Agitation Sedation Scale; SAM - Serratus anterior muscle; SPB - serratus plane block; TEB - thoracic epidural block; TPVB - thoracic paravertebral block.

References

1. Torre PA de la, Jones JW, Álvarez SL, et al. Axillary local anesthetic spread after the thoracic interfascial ultrasound block – a cadaveric and radiological evaluation. *Rev Bras Anesthesiol.* 2017;67:555-64.
2. Blanco R, Parras T, McDonnell JG, Prats-Galino A. Serratus plane block: A novel ultrasound-guided thoracic wall nerve block. *Anaesthesia.* 2013;68:1107-13.
3. De Cassai A, Bonvicini D, Ruol M, Correale C, Furnari M. Erector spinae plane block combined with a novel technique for selective brachial plexus block in breast cancer surgery. *Korean J Anesthesiol.* 2019;72:270-74.
4. Khetarpal R, Chatrath V, Dhawan A, Attri J. Regional anesthesia in difficult airway: The quest for a solution continues. *Anesth Essays Res.* 2016;10:178-83.
5. Sanllorente-Sebastián R, Rodríguez-Joris E, Avello-Taboada R, Fernández-López L, Ayerza-Casas V, Robador-Martínez D. Addition of serratus-intercostal plane block/BRILMA for arteriovenous access surgery. *Rev Esp Anesthesiol Reanim.* 2020;67:343-6.

Figure 1 A, Sonoanatomy of the serratus plane block (LDM, Latissimus Dorsi Muscle; SAM, Serratus Anterior Muscle). B, Image after surgical debridement. C, Sonoanatomy of supraclavicular brachial plexus block (BP, Brachial Plexus; SCA, Subclavian Artery).



Table 1 Axillary anatomy.

Walls	Boundaries	Inervation
Medial	External surface of 4 first ribs	Intercostal nerves
	Serratus anterior muscle	Long thoracic Nerve C5–C7
Posterior	Subscapularis muscle	Lower and upper subscapular nerves C5–C6
	Teres major muscle	Subscapular nerves
	Latissimus dorsi muscle tendon	Thoracodorsal nerve C8–T1
Anterior	Pectoralis major muscle	Lateral pectoral nerve C5–C7 and medial pectoral nerve C8–T1
	Pectoralis minor muscle	
	Pectoral and clavipectoral fascias	
Base	Axillary fascia, adipose layer, and skin	Intercostobrachial nerve T2–T3
		Medial brachial cutaneous nerve C8–T1
		Lateral cutaneous branches of intercostal nerves T3–T9
Apex	Cervicoaxillary canal	
Lateral	Space between the humerus and the insertions of the muscles of the anterior and posterior walls	

Source: Moore KL, Dalley AF, Agur AM. Clinically Oriented Anatomy Seventh Edition. Lippincott Williams & Wilkins; 2014. p. 851-870.