

Journal Pre-proof

A difficult airway approach in a merosin-deficient congenital muscular dystrophy patient – a case report

Jorge Pelicano Paulos, Vanessa Artilheiro, Catarina Cruz, Ana Pinto Carneiro



PII: S0104-0014(21)00149-4

DOI: <https://doi.org/10.1016/j.bjane.2021.03.018>

Reference: BJANE 744125

To appear in: *Brazilian Journal of Anesthesiology (English edition)*

Please cite this article as: { doi: <https://doi.org/>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

BJAN-D-20-00034 - Case Report**A difficult airway approach in a merosin-deficient congenital muscular dystrophy patient – a case report**

Jorge Pelicano Paulos^{a,*}, Vanessa Artilheiro^b, Catarina Cruz^c, Ana Pinto Carneiro^a

^a Centro Hospitalar Universitário Lisboa Central, Hospital Dona Estefânia, Anesthesiology Department, Lisbon, Portugal

^b Centro Hospitalar de Lisboa Ocidental, Anesthesiology Department, Lisbon, Portugal

^c Hospital de Cascais, Anesthesiology Department, Cascais, Portugal

*Corresponding author.

E-mail: pelicanopaulos@gmail.com (J.P. Paulos).

ORCID ID:

Jorge Pelicano Paulos - <https://orcid.org/0000-0001-7443-9874>

Received 4 May 2020; accepted 19 March 2021

Abstract

Merosin-deficient muscular dystrophy is caused by an autosomal recessive mutation on laminin- α 2 gene characterized by severe progressive muscle weakness associated with neuromuscular scoliosis and restrictive lung disease. In this case report, we describe an alternative airway approach performed in a child with anticipated difficult airway and merosin-deficient muscular dystrophy. Significant anesthetic implications may increase the perioperative risk, requiring accurate knowledge to anticipate an adequate management and provide patient-safety strategies.

KEYWORDS: Pediatrics; Orthopedics; Airway management; Intravenous anesthesia; Muscular dystrophy congenital, merosin negative

Introduction

Congenital muscular dystrophies (CMDs) are autosomal recessive disorders with compromised synthesis and regeneration of the muscle contractile proteins. Besides all

being rare and progressive, they differ in their clinical manifestations and genetic background.[1]

Merosin-deficient CMD (MD-CMD) has a prevalence of about 0,7/100.000 and is caused by a mutation on chromosome 6, in laminin- α 2 gene (or merosin) present in skeletal muscle fibers.[2] It accounts for nearly 40% of all CMD cases, the most common and severe form. Regardless, few reports of its anesthetic management were published in the literature. This case report contributes to the current knowledge related to airway management in MD-CMD patients, sustaining a case-oriented approach to an anticipated difficult airway.

Case report

A nine-year-old boy, ASA III, diagnosed with MD-CMD during his first months of life, presented a severe dorsolumbar scoliosis and an inherent restrictive lung disease with diminished lung volumes and frequent respiratory infections. A previous scoliosis correction surgery already showed that only videolaryngoscopy provided a successful airway approach due to Cormack-Lehane score IV on conventional laryngoscopy. Therefore, the child was now proposed for scoliosis definitive surgical treatment under intraoperative neurophysiology monitoring.

During preoperative assessment, the patient demonstrated reduced neck mobility, slight retrognathism and a persistent left flexion of the head, macroglossia and Mallampati III, predictors consistent with MD-CMD and a difficult airway approach as previously identified (Fig. 1).

Intraoperatively, an inhaled induction with sevoflurane ensuring spontaneous ventilation, without opioid adjuvants, was maintained until intubation, allowing an adequate level of unconsciousness and also a safe and dynamic airway assessment. Initially, McCoy laryngoscopy evidenced a Cormack-Lehane IV, confirming the difficult airway assessment. Secondly, McGrath videolaryngoscopy showed an anterior glottis and intubation could not be productively accomplished even with a tube introducer. To improve the airway approach success, McGrath videolaryngoscopy was combined with an optic fiberscope as a guide for intubation, which was successfully achieved with a 6.0-cuffed tube (Fig. 2). The entire intubation process lasted 20 minutes.

Maintenance with remifentanyl and propofol total intravenous anesthesia without neuromuscular blockade was performed avoiding any interference with evoked potential

acquisition. The surgical procedure was uneventful and the patient remained intubated in the Pediatric Intensive Care Unit for 24 hours. Extubation was planned after weaning of sedation and mechanical ventilation under Anesthesiology team supervision. No further complications occurred during postoperative period.

Discussion

It is widely recognized that MD-CMD patients typically require orthopedic procedures early in life, namely scoliosis correction; however, anesthetic management is particularly challenging and burdened with perioperative complications.[3] Before the procedure, an experienced multidisciplinary team should be involved to implement patient-safety strategies and provide an integrated care. This should include a comprehensive evaluation of the patient's global functional condition and anticipation of difficult airway predictors regarding their poor head control, marked neck and trunk hypotonia and limited mouth opening[1] as well as the increased risk of rhabdomyolysis.[3]

In what difficult airway concerns, fiberoptic intubation is the gold-standard, although it does not always provide a successful approach and no single airway device or technique will adapt in every clinical situation.[4] The option of a videolaryngoscopy approach after a Cormack-Lehane IV identification was an initial attempt to optimize glottis visualization as part of the process of difficult airway algorithm. Although the absence attempt of a flexible fibroscopy alone was an eventual limitation, we believe that a simultaneous approach of these two airway devices provided a better visualization through the videolaryngoscope screen and an easier control of the tube tip with the fiberscope in order to secure the airway. As stated by other authors,[5] this combined technique is advantageous to decrease tissue trauma particularly in patients with high Cormack-Lehane scores. Therefore, maintenance of spontaneous ventilation combined with a videolaryngoscopy-assisted fiberoptic intubation should be considered as an alternative to difficult but non-emergent airway scenarios in Pediatric Anesthesiology, specially when there is an association of a higher Cormack-Lehane score with other clinical difficult airway predictors.

Concerning timing of tracheal extubation, most authors suggest performing it awake after a complete reversal of neuromuscular blockade.[3] Non-invasive ventilation should be considered as a resource for immediate postoperative period, bearing in mind the possibility of respiratory failure and need of Intensive Care Unit admission.

Standard ASA monitoring must always be used and neuromuscular blockade stimulation is recommended mostly in cases of severe hypotonia. Albeit MD-CMD patients are not prone to malignant hyperthermia, its theoretical risk still exists as a muscular dystrophy, along with anesthesia-induced rhabdomyolysis. Both may induce arrhythmias and respiratory failure exacerbation. Hence, general recommendations include a trigger-free anesthetic technique and prefer an inhaled induction as a less traumatic experience. Multimodal analgesia seems especially important to reduce the inherent risk of most complications.

Learning points

Muscular dystrophies share significant perioperative risk, namely a difficult airway approach or MH. A combined airway approach involving a McGrath videolaryngoscopy and a flexible fiberscope provides a valid alternative approach of an expected difficult airway in a child with MD-CMD. A multidisciplinary strategy guarantees the most effective perioperative care, minimizing any issues related to the disease and anesthetic management.

Conflicts of interest

The authors declare no conflict of interest.

Acknowledgments

We would like to thank Dr Teresa Cenicante, MD for all her availability and support on reporting and working in case reports such as this one as the responsible of Pediatric Anesthesiology in Centro Hospitalar Universitário Lisboa Central.

References

- [1] Jimenez N, Song K, Lynn AM. Hemodynamic instability during prone spine surgery in a patient with merosin-deficient congenital muscular dystrophy. *Paediatr Anaesth.* 2013;23:294–6.
- [2] Pregardien C, Pirotte T, Veyckemans F. Pressure-support ventilation in a child with merosin-deficient congenital muscular dystrophy under sevoflurane anesthesia. *Acta*

Anaesthesiol Belg. 2016;67:139–41.

[3] Echeverry-Marín PC, Bustamante-Vega ÁM. Anesthetic implications of muscular dystrophies. Rev Colomb Anesthesiol. 2018;46:228-39.

[4] Matic AA. Use of the Airtraq® with a fiberoptic bronchoscope in a difficult intubation outside the operating room. Can J Anesth. 2008;55:561-2.

[5] Sgalambro F, Sanfilippo F, Santonocito C, Caltavuturo C, Grillo C. Virtual laryngoscopy and combined laryngoscopic-bronchoscopic approach for safe management of obstructive upper airways lesions. Br J Anaesth. 2014;113:304-6.

Figure 1 – Preoperative evaluation of the airway.



Figure 2 – Status after airway management.

