

Journal Pre-proof

A simple technique to maintain intraoperative head and neck neutrality

Anthony M.-H. Ho , Gregory Klar , Glenio Bitencourt Mizubuti

PII: S0104-0014(21)00403-6
DOI: <https://doi.org/10.1016/j.bjane.2021.11.006>
Reference: BJANE 744327



To appear in: *Brazilian Journal of Anesthesiology (English edition)*

Received date: 16 September 2021
Accepted date: 13 November 2021

Please cite this article as: Anthony M.-H. Ho , Gregory Klar , Glenio Bitencourt Mizubuti , A simple technique to maintain intraoperative head and neck neutrality, *Brazilian Journal of Anesthesiology (English edition)* (2021), doi: <https://doi.org/10.1016/j.bjane.2021.11.006>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Editora Ltda. on behalf of Sociedade Brasileira de Anestesiologia. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

BJAN-D-21-00412 - Clinical Image

A simple technique to maintain intraoperative head and neck neutrality

Anthony M.-H. Ho, Gregory Klar, Glenio Bitencourt Mizubuti*

Queen's University, Department of Anesthesiology and Perioperative Medicine,
Kingston, Canada

*** Corresponding author.**

E-mail: Gleniomizubuti@hotmail.com (G.B. Mizubuti).

ORCID ID:

Anthony M.-H. Ho - 0000-0002-9721-4499

Gregory Klar - 0001-9166-0108

Glenio B. Mizubuti - 0000-0002-5515-829X

Received 16 September 2021; accepted 13 November 2021

Keywords

Patient safety; Intraoperative head position; Intraoperative neck position; Laparoscopic cholecystectomy; Patient positioning; Intraoperative injury

During laparoscopic cholecystectomy, the left-tilt plus Trendelenburg positioning required can result in the head and neck being turned to an extreme left and/or flexed position (Fig. 1A). In certain upper extremity surgeries, the patient is positioned at the edge of the table and the patient's head is at risk of drifting unnoticed off the edge (Fig. 1B).

Excessive head rotation is potentially harmful. Weintraub and Khoury have shown that major occlusion of the vertebral artery occurs in 56.4% and 13.3% of

patients with (mean age 71 years) and without (mean age 66 years) vascular risk factors, respectively, upon maximum head rotation.[1] In adults, Burbridge et al have found that 80% of the internal jugular vein is occluded during head turn of $\sim 55^\circ$, potentially raising intracranial pressure.[2] Carotid dissection has been reported in a 36-year-old healthy woman clutching her phone between her head and shoulder for 32 minutes.[3] One may speculate that carotid compression plus reverse Trendelenburg positioning could compromise cerebral perfusion. Patients with cervical spine pathology may not tolerate head flexion and tilt. Stretching of the contralateral ligaments, muscles, and nerves create strain.

Figure 1C shows how use of the anesthesia circuit holder can maintain the patient's head in a neutral position. Figure 1D shows the same arrangement reinforced to stabilize the head and neck. Finally, the anesthesia circuit holder is shown in Figure 1E.

Conflicts of interest

The authors declare no conflicts of interest.

References

1. Weintraub MI, Khoury A. Critical neck position as an independent risk factor for posterior circulation stroke. A magnetic resonance angiographic analysis. *J Neuroimag.* 1995;5:16-22.
2. Burbridge MA, Min JG, Jaffe RA. Effect of head rotation on jugular vein patency under general anesthesia. *Can J Neurol Sci.* 2019;46:355-7.
3. Mourad J-J, Girerd X, Safar M. Carotid-artery dissection after a prolonged telephone call. *N Engl J Med.* 1977;336:516.

Figure 1 - Typical head positions during surgery. (A) Significant flexion and left lateral neck/head tilt during typical patient positioning for laparoscopic cholecystectomy (i.e., reverse Trendelenburg with left lateral tilt of the surgical table). In this example, the (optional) pillow/cushion commonly used to support the patient's head is absent. (B) Patient positioned at the edge of the surgical table as commonly observed in upper extremity surgeries. Notably, the surgical table remains in neutral position, but the patient's head is at risk of falling off to the side of the surgical table (especially with surgical manipulation) despite the use of a pillow/cushion to support it. Finally, panels

(C) and (D) depict the simple use of the anesthesia circuit holder (E) to maintain the patient's head/neck in neutral position during cholecystectomy (reverse Trendelenburg and steep left lateral tilt of the surgical table) and upper extremity surgery (surgical table in neutral position). Note the use of a clamp (green arrow) to secure the circuit holder against the mattress and/or its covering sheet to prevent it from slipping along with the patient's head, thereby further stabilizing the head/neck position. Also, a towel (red arrows) is used for providing comfort and to isolate the patient's face/head from the metallic holder to prevent against electrocautery-induced skin burns.

